Automated Stratigraphic Mapping using Convolutional Neural Networks

A. M. Annex*, K. W. Lewis



Mapping Layered Deposits on Mars


Layered deposits have been found all over Mars.

We don't know why/how!

We make high resolution DEMs (elevation models) with HiRISE images

We measure <u>orientations</u> and <u>layer</u> thickness to examine spatial trends in the deposits. (Annex et al. 2018 in prep)

It is a lot of work!



#48 A. Annex




Training data & examples


Human labels (yellow lines), 1 meter per pixel images


- 1. Improved labels with VGG Image Annotator (VIA)
- 2. Training data covers diverse morphologies of layered deposits.
- 3. 797 images total
- 4. Overall, an imbalanced classification problem!


Used modified U-net architecture (added batchnorm). keras data augmentation





True Positives













True Negatives





U-net metrics




input 0 label 0 prediction epoch: 2 50 50 50 100 100 -100 -150 -150 150 200 -200 200 250 -250 100 50 100 150 250 150 200 250 150 ROC curve, AUC: 0.912 Precision-Recall curve: AP=0.428

Human "ground truth"


Computer Prediction


HiRISE Input

| T=382564 F=10627484 | False (P) | True (P) |
|------------------------|-----------|----------|
| False (A) | 0.964 | 0.035 |
| True (A) | 0.261 | 0.738 |




#48 A. Annex






Results from CNN mapping

- ~40k strata mapped
- 16k layer thicknesses
- 5k after naive filtering
- Human time: several days, months in human "wall time"
- Computer time: a few hours to train, minutes to run predicts & fits
- Need to work on better post processing and filtering...





Conclusions

- 1. Planetary Sciences can benefit from dl/ml to automate tasks that are not science! Save grad students from scientific busy work.
- 2. U-net produced good results, with multiple morphologies of layered units present in training data.
- 3. Newer architectures to test: 100 layer Tiramisu, GANs.
- 4. Need to try older methods: HOG+SVM.
- 5. Need to use CV to compare these models.
- 6. Mutual benefits between EO and Planetary fields.







